

SHARDA METALDEHYDE 3GB

SHARDA SNAIL AND SLUG BAIT



Reg. No. L 9520, Act No. 36 of 1947

A compressed pellet bait for the control of snails
and slugs in crops and the home garden.
Ready for use.

Reg. Nr. L 9520, Wet Nr. 36 van 1947

'n Saamgepersde pilletjie lokaas vir die beheer
van slakke in gewasse en die huistuin.
Gereed om gebruik te word.

Active ingredient
Metaldehyde

30 g/kg

Aktiewe bestanddeel
Metaldehied

Manufactured and Registered by Vervaardig en Geregistreer deur
Sharda International Africa cc
Company Reg. No. 2010/002268/07 Maatskappy Reg. Nr.
P O Box/Posbus 82021, Southdale 2135.
Tel: 031-266-3264

Date of Manufacture
Batch number

Datum van vervaardiging
Lot nommer

20Kg



CAUTION / VERSIGTIG

SHARDA METALDEHYDE 3GB

WARNINGS

- Do not apply directly onto edible crops. Apply between plants.
- Poisonous if swallowed.
- Keep out of the reach of children, uninformed persons, poultry and animals.
- Store in a cool, dry place away from food and feedstuffs.

PRECAUTIONS

- Prevent contamination of food, feed, water sources and fish ponds.
- Wash hands after handling the pellets.
- Destroy the empty container and do not re-use for any other purpose.

DIRECTIONS FOR USE

Use only as Directed

- Scatter the pellets evenly on moist soil. Do not heap pellets in a pile.
- Apply late afternoon/early evening.
- Do not apply to the edible portion of any crop.
- Apply the higher rate for high infestations.

WAARSKUWINGS

- Moet nie direk op eetbare gewasse toedien nie. Dien toe tussen plante.
- Giftig indien ingestuk.
- Hou buite die bereik van kinders, oningeligte persone, pluimvee en diere.
- Stoor in 'n koel, droë plek weg van voedsel en voer.

VOORSORGMATREËLS

- Voorkom besoedeling van voedsel, voer, waterbronne en visdamme.
- Was hande na hantering van die pilletjies.
- Vernietig die leë houer en moet nie hergebruik vir enige ander doel nie.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

- Strooi die pilletjies egalig op klam grond. Moet nie pilletjies ophoop nie.
- Dien toe in die laat middag/vroeë aand.
- Moet nie op die eetbare gedeelte van 'n gewas toedien nie.
- Gebruik die hoër dosis in geval van hoë infestasië.

SITUATION	DOSAGE	REMARKS
Home garden	8 – 10 g/10 m ² (High infestation: 20 – 30 g/10 m ²)	Apply preventatively when the first snails or slugs are noticed. Repeat as necessary.
Crops, Flowers, Vegetables.	8 – 10 kg/ha (High infestation: 20 – 30 kg/ha)	Apply preventatively before the first snails or slugs are noticed and cause damage. Repeat as necessary.
Bananas	8 – 10 kg/ha (High infestation: 20 – 30 kg/ha)	Apply around the base of each plant. Apply preventatively before the first snails or slugs are noticed and cause damage. Repeat at 3 – 4 week intervals as necessary.
Grapes	8 – 10 kg/ha (High infestation: 20 – 30 kg/ha)	Apply around the stems of plants and on the ridges. Apply preventatively before budding and the first snails or slugs are noticed and cause damage. Repeat at 3 week intervals or as necessary.

SITUASIE	DOSIS	OPMERKINGS
Huistuin	8 – 10 g/10 m ² (Hoë infestasië: 20 – 30 g/10 m ²)	Behandel voorkomend sodra die eerste slakke opgemerk word. Herhaal soos benodig.
Gewasse, Blomme, Groente	8 – 10 kg/ha (Hoë infestasië: 20 – 30 kg/ha)	Behandel voorkomend sodra die eerste slakke opgemerk word. Herhaal soos benodig.
Piesangs	8 – 10 kg/ha (Hoë infestasië: 20 – 30 kg/ha)	Strooi om elke plant. Behandel voorkomend sodra die eerste slakke opgemerk word en skade veroorsaak. Herhaal elke 3 – 4 weke of soos benodig.
Druiewe	8 – 10 kg/ha (Hoë infestasië: 20 – 30 kg/ha)	Behandel om die stokke en op die bankies. Behandel voorkomend voor die bot groei stadium bereik word of sodra die eerste slakke opgemerk word en skade veroorsaak. Herhaal met 3 week tussenposes of soos benodig.

