



AZOX FORTE 50 WG

Reg. No. L10486, Act No. 36 of 1947 | Reg. Nr. L10486, Wet Nr. 36 van 1947

Read the label before use
Keep out of reach of children and animals

Azox Forte 50 WG is a water dispersible granule systemic and translaminar fungicide for the control of powdery mildew and downy mildew in wine and table grapes.

Azox Forte 50 WG is 'n waterdispergeerbare korrel sistemiese en translaminêre swamdoder vir die beheer van poeieragtige skimmel en donsige skimmel in wyn- en tafeldruive.

Warning
Hazard Statement

Harmful if inhaled.

Very toxic to aquatic life with long lasting effects.

Precautionary Statements

Avoid breathing dust/fume/gas/mist/
vapours/spray.

Use only outdoors or in a well-ventilated area



FUNGICIDE GROUP 11

ACTIVE INGREDIENTS:

Azoxystrobin 500 g/kg

Registered by | Geregistreer deur

Sharda International Africa (PTY) LTD

Reg. No./Reg. Nr. 2010/002268/07

P. O. Box/Posbus 82021,

Southdale, 2135

Tel: 031-764 3011

SWAMDODER GROEP 11

AKTIEWE BESTANDDELE:

Asoksiestrobin 500 g/kg

BATCH NO.		LOT NR.	
DATE OF MANUFACTURE		DATUM VAN VERVAARDIGER	
EXPIRY DATE		VERVAL DATUM	

NET CONTENTS /
NETTO INHOUD

1 Kg

UN Number 3077

National Poison helpline number/ emergency contact 086 155 5171

WARNINGS:

- **Withholding periods:** Minimum number of days between last application and harvest.
- **Wine and table grapes:** **14 days**

NOTE: Compliance with these withholding periods will ensure that residues do not exceed local Maximum Residue Limits, but the import tolerances of other countries might possibly be exceeded. If the crop to be treated is intended for export, consult the relevant importer or exporting body regarding the use of this product. Maximum Residue Limits and recommended withholding periods.

- Harmful if inhaled.
- Harmful if ingested.
- Very toxic to aquatic life with long lasting effects.
- Avoid contact with skin or eyes since product may cause eye and skin irritation.
- Store away from food and feed.
- Keep out of reach of children, uninformed persons and animals.
- **Re-entry:** Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic, and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Do not eat, drink or smoke whilst handling this product.
- Avoid breathing dust/fume/gas/mist/ vapours/spray.
- Use only outdoors or in a well-ventilated area
- Avoid release to the environment.
- IF INHALED: Remove person to fresh air and keep comfortable for breathing.
- Collect spillage.
- Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Wear rubber gloves and face shield when handling the concentrate. If you get it in your eyes, flush it out at once. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them.
- Containers and packages must be completely emptied before being disposed. Shake out thoroughly into the mixing or spray tank before destroying the empty container.
- While spraying, avoid contact with the spray as much as you can. Avoid spray drift onto other crops, grazing, rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams. Destroy empty container and do not use for any other purpose. Change and wash your work clothes. Wash yourself.

RELEVANT SUBSTANCES:

Components	Concentration % by weight
Azoxystrobin	< 50

RESISTANCE MANAGEMENT:

For resistance management, **Azox Forte 50 WG** is a group code 11 fungicide. Any fungus population may contain individuals naturally resistant to **Azox Forte 50 WG** and other group code 11 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **Azox Forte 50 WG** or any other group code 11 fungicides.

To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes.
- Refer to individual product labels when alternating products or when using tank mixtures with products in fungicide group codes 7, 27, 28, 29, 30 and M.
- Integrate other control methods (chemical, cultural, biological) into disease control programs.

Azox Forte 50 WG should always be applied as a preventive treatment. Do not apply less than two, or more than three, consecutive sprays per season.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.

Poisons Helpline: 086 155 5171

Mixing instructions:

Mix the required amount of **Azox Forte 50 WG** with a small quantity of water to form a smooth cream without any remaining granules. Half fill the spray tank with water. Add the creamed product to the tank through a sieve. Top up to the required volume while agitating continuously. Maintain agitation throughout application and do not allow tank mixture to stand still for any period of time.

CROP / DISEASE	DOSAGE
WINE AND TABLE GRAPES Downy mildew <i>(Plasmopara viticola)</i> and Powdery mildew <i>(Oidium)</i> <i>(Uncinula necator)</i>	10 – 14 day programme: 40 g / 100 ℓ water
REMARKS: High volume application: Commence application when the longest shoots are ± 10 cm long and repeat at 10 – 14 day intervals for not more than 3 consecutive application per season. For the control of powdery mildew, a spray programme must already be started at 2,5 cm shoot length with remedies from a different chemical class. Wine grapes: Pre blossom: apply in 250 – 500 ℓ spray mixture / ha. During blossom: apply in 500 – 750 ℓ spray mixture / ha. Post- blossom: apply in 750 – 1000 ℓ spray mixture / ha. Table grapes: Pre-blossom: apply in 500 – 750 ℓ spray mixture / ha. During blossom: apply in 750 – 1000 ℓ spray mixture / ha. Post-blossom: apply in 1000 – 1500 ℓ spray mixture / ha.	
NOTE: Azox Forte 50 WG can be applied at any stage during the spray programme but the most suitable time is during the blossom to pea berry stage.	
IMPORTANT: 1. Azox Forte 50 WG must always be applied as a preventive treatment. 2. Do not apply less than two, or more than three, consecutive sprays of Azox Forte 50 WG per season. 3. After the last Azox Forte 50 WG application, change to a fungicide from a different chemical class e.g. HEXACONAZOLE 50 g / ℓ. (Consult the HEXACONAZOLE 50 g / ℓ label for full particulars. 4. In cases where both diseases are present after Azox Forte 50 WG has already been applied, use other registered products which do not belong to the same chemical class as Azox Forte 50 WG to control the diseases. 5. Do not apply other strobilurin containing fungicides e.g. TRIFLOXYSTROBIN 500 g/kg WG and KRESOXIM-METHYL 500g/kg WG or any other fungicides which have the same mode of action, in the same spray programme with Azox Forte 50 WG 6. Do not apply Azox Forte 50 WG as the first or final application in the spray programme. 7. Do not apply Azox Forte 50 WG at reduced rates. 8. Do not apply Azox Forte 50 WG to nursery grape vine cuttings.	