



VOLT 80 WG

Reg. No. L10465, Act No. 36 of 1947 | Reg. Nr. L10465, Wet Nr. 36 van 1947

Read the label before use
Keep out of reach of children and animals

Volt 80 WG is a water dispersible granule organic based fungicide for the preventive control of anthracnose, downy mildew, dead arm disease and Botrytis rot in wine grapes, dead arm disease, anthracnose and downy mildew pre-blossom in table grapes as well as late and early blight in potatoes and tomatoes.

Hazard Statements

Toxic if inhaled
Causes serious eye irritation
May cause an allergic reaction
Suspected of causing cancer
Toxic to aquatic life with long lasting effects

Precautionary Statements

Avoid breathing mist / spray
Wear protective gloves
Avoid release to the environment.

FUNGICIDE GROUP CODE Y

ACTIVE INGREDIENT:

Folpet (phtalimide) 800 g/kg

Manufactured for and registered by/Vervaardig
vir en geregistreer deur:

Sharda International Africa (PTY) LTD

Reg. No./Reg. Nr. 2010/002268/07

P. O. Box/Posbus 82021, Southdale, 2135

Tel: 031-764 3011

Volt 80 WG is 'n waterdispergeerbare korrel organies-gebaseerde swamdoder vir die voorkomende beheer van antraknose, donsige skimmel, streepvleksierte en Botritis-vrot in wyndruiwe; streepvleksierte, antraknose en donsige skimmel voor-blom in tafeldruiwe asook laat- en vroeëroes in aartappels en tamaties.



Danger

SWAMDODERGROEP CODE Y

AKTIEWE BESTANDDEEL:

Folpet (ptalimied) 800 g/kg

BATCH NUMBER
LOT NOMMER

DATE OF EXPIRY
VERVAL DATUM

NET CONTENTS /
NETTO INHOUD

25 Kg

UN No. / VN Nr. 3077

National Poison helpline number/ emergency contact 086 155 5171

WARNINGS:

- **Withholding periods:** (Minimum number of days between last application and harvest).

Tomatoes3 days

Table grapesnot later than pre-blossom stage.

Wine grapes14 days

- Toxic if inhaled
- Causes serious eye damage
- May cause an allergic skin irritation
- Suspected of causing cancer
- Very toxic to aquatic life
- Store in a cool place away from food and feed.
- Keep out of reach of children, uninformed persons and animals.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC, AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE DISEASE TO THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHER DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

- Avoid breathing dust/fume/gas/mist/vapours/spray.
- Contaminated work clothing should not be allowed out of the workplace.
- Wear protective gloves/protective clothing/eye protection/face protection.
- Use only outdoors or in a well-ventilated area.
- Obtain special instructions before use.
- Do not handle until all safety precautions have been read and understood.
- Use personal protective equipment as required.
- Wash face, hands and any exposed skin thoroughly after handling.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF exposed or concerned: Get medical advice/attention.
- IF ON SKIN: Wash with plenty of soap and water.
- If skin irritation or rash occurs: Get medical advice/attention.
- Specific treatment (see supplemental instructions on the administration of antidotes on this label).
- Wash contaminated clothing before reuse.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- If eye irritation persists: Get medical advice/attention.
- IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.
- Call a POISON CENTER or doctor/physician if you feel unwell.
- Avoid release to the environment.
- Collect spillage
- Do not eat, drink or smoke whilst handling this product.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Containers and packages must be completely emptied before being disposed of. Shake out thoroughly into the mixing or spray tank before destroying the empty container.
- While spraying, avoid contact with the spray as much as you can. Avoid spray drift onto other crops, grazing rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.
- Destroy empty container by perforating and flattening and do not use for any other purpose.

RELEVANT SUBSTANCES:

Components	%
Folpet	< 85%
Sodium alkyl-naphthalene sulfonate, formaldehyde condensate proprietary	<5%
Alkyl-naphthalene-sulphonic acid	<2 %

DIRECTIONS OF USE: Use only as indicated.

Compatibility:

Volt 80 WG is compatible with most insecticides and fungicides commonly used with the exception of Lime, Lime sulphur and Bordeaux mixture.

Mixing and application instructions:

Mix the required quantity of **Volt 80 WG** separately in a small amount of water to make a smooth cream before adding it to the recommended amount of water in the spray tank. Start agitating whilst adding **Volt 80 WG** into the spray tank and maintain agitation throughout the application operation. Spray thoroughly to ensure complete coverage of the plant, including the under-surface of the leaves. **Volt 80 WG** is recommended as a preventive treatment.

CROP AND DISEASE	DOSAGE	REMARKS
Wine and table grapes Anthracnose, Deadarm disease and Botrytis rot	125 g / 100 ℓ water	Apply every two weeks or weekly when weather conditions are favourable for the development of the disease.
Downy mildew	125 g / 100 ℓ water	Pre-blossom: Apply the first application when shoots are 10 cm long. Repeat at 7 – 10 day intervals depending on weather conditions which will favour the development of the disease. Blossom to 80 % petal drop: Only apply when the flowering period lasts for more than two weeks. Post-blossom: Apply twice or more at 7 – 14 day intervals, depending on weather conditions.
Potatoes Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>)	1.25 kg / ha	Preventive treatment: Apply from emergence at 7 – 10 day intervals in not less than 500 ℓ water / ha in the pre-blossom stage in 1000 ℓ water / ha in the post blossom stage. For the control of early and late blight, particularly after flowering, it is important to obtain thorough coverage. Add a suitable adjuvant at the recommended rate.
Tomatoes Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>)	1.25 kg / ha	Apply as a preventive application shortly after transplant and before symptoms of the disease appear. Depending on plant size apply in 300 – 1500 ℓ water / ha and ensure thorough coverage of both upper and lower leaf surfaces. Repeat application at 7 – 10 day intervals.