



AZOX FORTE 50 WG

Reg No. .10486 | Act no/Wet Nr 36 of/van1947

Azox Forte 50 WG is a water dispersible granule systemic and translaminar fungicide for the control of powdery mildew and downy mildew in wine and table grapes.

Azox Forte 50 WG is 'n waterdispergeerbare korrel sistemiese en translaminêre swamdoder vir die beheer van poeieragtige skimmel en donsige skimmel in wyn- en tafeldruive.

FUNGICIDE GROUP 11

ACTIVE CONSTITUENTS:

Azoxystrobin 500 g/kg

Registration Holder / Registrasiehouer:

Sharda International Africa (Pty) Ltd
Company Reg. No. 2010/002268/07
P O Box 82021, Southdale 2135.
Tel: 0317643011/0878222397

SWAMDODER GROEP 11

AKTIEWE BESTANDDELE:

Asoksiestrobin 500 g/kg

BATCH NO/LOT NR	
Expiry Date Verval Datum	

NET CONTENTS /
NETTO INHOUD

1Kg

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE OR OPENING
BELANGRIK: LEES AANGEHEGTE PAMFLET VOOR GEBRUIK OF OOPMAAK



CAUTION
VERSIGTIG



WARNINGS:

- **Withholding periods:** Minimum number of days between last application and harvest.

Wine and table grapes: 14 days

NOTE: Compliance with these withholding periods will ensure that residues do not exceed local Maximum Residue Limits, but the import tolerances of other countries might possibly be exceeded. If the crop to be treated is intended for export, consult the relevant importer or exporting body regarding the use of this product. Maximum Residue Limits and recommended withholding periods.

- Harmful if ingested or swallowed.
- Toxic to fish and harmful to other aquatic life.
- Avoid contact with skin or eyes since product may cause eye and skin irritation.
- Store away from food and feed.
- Keep out of reach of children, uninformed persons and animals.
- **Re-entry:** Do not enter treated area until spray deposit has dried unless wearing protective clothing.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC, AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURENCE OF RESISTANCE OF THE DISEASE TO THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHER DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

- Do not eat, drink or smoke whilst handling this product.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Wear rubber gloves and face shield when handling the concentrate. If you get it in your eyes, flush it out at once. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them.
- Containers and packages must be completely emptied before being disposed. Shake out thoroughly into the mixing or spray tank before destroying the empty container.
- While spraying, avoid contact with the spray as much as you can. Avoid spray drift onto other crops, grazing, rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams. Destroy empty container and do not use for any other purpose. Change and wash your work clothes. Wash yourself.

RESISTANCE MANAGEMENT:

For resistance management, **Azox Forte 50 WG** is a group code 11 fungicide. Any fungus population may contain individuals naturally resistant to **Azox Forte 50 WG** and other group code 11 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **Azox Forte 50 WG** or any other group code 11 fungicides.

To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes.
- Refer to individual product labels when alternating products or when using tank mixtures with products in fungicide group codes 7, 27, 28, 29, 30 and M.
- Integrate other control methods (chemical, cultural, biological) into disease control programs.

Azox Forte 50 WG should always be applied as a preventive treatment. Do not apply less than two, or more than three, consecutive sprays per season.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.

Poisons Helpline: 0861 555 777

Mixing instructions:

Mix the required amount of **Azox Forte 50 WG** with a small quantity of water to form a smooth cream without any remaining granules. Half fill the spray tank with water. Add the creamed product to the tank through a sieve. Top up to the required volume while agitating continuously. Maintain agitation throughout application and do not allow tank mixture to stand still for any period of time.

CROP / DISEASE	DOSAGE
WINE AND TABLE GRAPES Downy mildew (<i>Plasmopara viticola</i>) and Powdery mildew (<i>Oidium</i>) (<i>Uncinula necator</i>)	10 – 14 day programme: 40 g / 100 ℓ water
REMARKS: High volume application: Commence application when the longest shoots are ± 10 cm long and repeat at 10 – 14 day intervals for not more than 3 consecutive application per season. For the control of powdery mildew, a spray programme must already be started at 2,5 cm shoot length with remedies from a different chemical class.	
Wine grapes: Pre blossom: apply in 250 – 500 ℓ spray mixture / ha. During blossom: apply in 500 – 750 ℓ spray mixture / ha. Post- blossom: apply in 750 – 1000 ℓ spray mixture / ha.	
Table grapes: Pre-blossom: apply in 500 – 750 ℓ spray mixture / ha. During blossom: apply in 750 – 1000 ℓ spray mixture / ha. Post-blossom: apply in 1000 – 1500 ℓ spray mixture / ha.	
NOTE: Azox Forte 50 WG can be applied at any stage during the spray programme but the most suitable time is during the blossom to pea berry stage.	
IMPORTANT: 1. Azox Forte 50 WG must always be applied as a preventive treatment. 2. Do not apply less than two, or more than three, consecutive sprays of Azox Forte 50 WG per season. 3. After the last Azox Forte 50 WG application, change to a fungicide from a different chemical class e.g. HEXACONAZOLE 50 g / ℓ. (Consult the HEXACONAZOLE 50 g / ℓ label for full particulars). 4. In cases where both diseases are present after Azox Forte 50 WG has already been applied, use other registered products which do not belong to the same chemical class as Azox Forte 50 WG to control the diseases. 5. Do not apply other strobilurin containing fungicides e.g. TRIFLOXYSTROBIN 500 g / kg and KRESOXIM-METHYL 500 g / kg or any other fungicides which have the same mode of action, in the same spray programme with Azox Forte 50 WG . 6. Do not apply Azox Forte 50 WG as the first or final application in the spray programme. 7. Do not apply Azox Forte 50 WG at reduced rates. 8. Do not apply Azox Forte 50 WG to nursery grape vine cuttings.	

WAARSKUWINGS:

- **Onthoudingsperiodes:** Minimum aantal dae tussen laaste toediening en oes.

Wyn- en tafeldruive:.....14 dae

LET WEL: Nakoming van hierdie onthoudingsperiodes sal verseker dat die plaaslike Maksimum Residu Limiete nie oorskry word nie, maar mag dalk nie voldoen aan die invoervereistes van ander lande nie. Indien die gewas wat behandel gaan word bestem is vir uitvoer, skakel die betrokke invoerder of uitvoerliggaam rakende die gebruik van hierdie produk, maksimum residu vereistes en aanbevole onthoudingsperiodes.

- Skadelik indien ingeneem, ingesluk of ingeasem word.
- Giftig vir visse en skadeling vir ander waterorganismes.
- Voorkom kontak met die oë en vel aangesien die middel oog- en velirritasie mag veroorsaak.
- Berg weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.
- **Herbtredding:** Moet nie behandelde gebiede betree alvorens die spuitneerslag droog is nie tensy beskermende oorklere gedra word.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GEOTOETS IS, WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE, AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE GROND-, KLIMAATS- EN BERGINGSTOESTANDE, KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE SIEKTE TEEN DIE BETROKKE MIDDEL ASOOK DIE METODE, TYD EN AKKURAAKHEID VAN TOEDIENING. VERDER AANVAAN DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR GEBREK AAN PRESTASIE VAN DIE BETROKKE MIDDEL AS GEVOLG VAN DIE VERSUM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMATREËLS:

- Moet nie eet, drink of rook terwyl die produk hanteer word nie.
- Vermyn kontaminasie van voedsel, voer, drinkwater en eetgerei.
- Dra rubberhandskoene en gesigskerm wanneer die konsentraat hanteer word. Indien dit in u oë kom, spoel onmiddellik uit. Indien dit op u vel kom, was dit dadelik af. Indien dit op u kleres mors, trek uit en was dit.
- Houers en verpakkings moet heeltemal leeg gemaak word voordat hulle vernietig word. Skud die houer deeglik uit in die meng- of spuittenk voordat leë houer vernietig word.
- Vermyn kontak met die spuitstof sover moontlik tydens toediening. Vermyn oorwaai van spuitnewel na ander gewasse, weiding, riviere of damme.
- Maak toediener deeglik skoon na gebruik en gooi waswater waar dit nie gewasse, weiding, riviere of damme sal besoedel nie. Vernietig leë houer en moet dit nie vir enige ander doel gebruik nie. Trek werksklere uit en was dit. Was u self.

WEERSTANDBESTUUR:

Azox Forte 50 WG is 'n groepkode 11 swamdoder. Enige populasie van 'n spesifieke swam mag individue insluit wat 'n natuurlike weerstand teen **Azox Forte 50 WG**, of enige ander groepkode 11 swamdoder het. Indien hierdie swamdoders in herhaaldelike en eksklusiewe programme aangewend word, kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **Azox Forte 50 WG** of enige ander groepkode 11 swamdoder, beheer word nie.

Om weerstand teen swamdoders te vertraag:

- Vermyn die herhaaldelike en eksklusiewe gebruik van swamdoders in dieselfde swamdoder groepkode. Wissel af met, of gebruik tenkmengsels van produkte in verskillende swamdoder groepcodes.
- Sien toepaslike individuele produketikette wanneer daar afgewissel word, of tenkmengsels gebruik word, met produkte in swamdoder groepcodes 7, 27, 28, 29, 30 of M.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in siektebeheerprogramme.

Azox Forte 50 WG moet altyd voorkomend toegedien word as deel van 'n siektebeheerprogram. Moenie minder as twee, of meer as drie toedienings per seisoen toedien nie.

Vir spesifieke inligting oor weerstandbestuur kontak die registrasiehouer van hierdie produk.

GBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

Vergiftigings hulplyn: 0861 555 777

Menginstruksies:

Meng die verlangde hoeveelheid **Azox Forte 50 WG** met 'n klein hoeveelheid water om 'n sagte room te vorm sonder enige oorblywende korrels. Maak die spuittenk halfvol met water. Voeg die room deur 'n sif in die tenk. Maak die tenk vol met water tot die verlangde volume terwyl aanhoudend geroer word. Hou aan om mengsel te roer tydens toediening. Moet nie tenkmengsel laat stilstaan nie, roer voortdurend.

GEWAS / SIEKTE	DOSIS
WYN- EN TAFELDRIUWE: Donsige skimmel (<i>Plasmopara viticola</i>) Poeieragtige skimmel (<i>Uncinula necator</i>)	10 - 14 dae programme: 40 g / 100 ℓ water
OPMERKINGS: Hoë volume toediening: Begin toediening wanneer die langste lote ± 10 cm lank is en herhaal met 10 – 14 dae tussenposes vir meer as 3 agtereenvolgende toedienings nie. Vir die beheer van poeieragtige skimmel, moet 'n programbespuiting reeds vanaf 2,5 cm lootlengte gevolg word, met middels uit 'n ander chemiese klas.	
Wyn-druive: Voor blom: dien toe in 250 – 500 ℓ spuitmengsel / ha. Tydens blom: dien toe in 500 – 750 ℓ spuitmengsel / ha. Na blom: dien toe in 750 – 1000 ℓ spuitmengsel / ha.	
Tafeldruive: Voor blom: dien toe in 500 – 750 ℓ spuitmengsel / ha. Tydens blom: dien toe in 750 – 1000 ℓ spuitmengsel / ha. Na blom: dien toe in 1000 – 1500 ℓ spuitmengsel / ha.	
LET WEL: Azox Forte 50 WG kan op enige stadium gedurende die spuitprogram toegedien word maar die mees geskikte tyd is gedurende die blom- tot erjie- korrelstadium.	
BELANGRIK: 1. Azox Forte 50 WG moet altyd as 'n voorkomende bespuiting toegedien word. 2. Moet nie minder as twee, of meer as 3 bespuitings gee van Azox Forte 50 WG per seisoen nie. 3. Na die laaste Azox Forte 50 WG toediening, verander na 'n swamdoder met 'n ander chemiese klas bv. HEXACONAZOOL. (Kyk na die HEXACONAZOOL etiket vir volle besonderhede). 4. In gevalle waar albei siektes teenwoordig is nadat Azox Forte 50 WG reeds toegedien is, gebruik ander swamdoders wat nie aan die selfde chemiese klas as Azox Forte 50 WG behoort nie om die siektes te beheer. 5. Moet nie ander strobilurienbevattende swamdoders bv. TRIFLOKSISTROBIEN 500 g/kg WG en KRESOXIM-METIEL 500 g/kg WG of enige ander swamdoders wat dieselfde wyse van werking het, saam met Azox Forte 50 WG in dieselfde spuitprogram toedien nie. 6. Moet nie Azox Forte 50 WG as die eerste of laaste toediening in die spuitprogram toedien nie. 7. Moet nie Azox Forte 50 WG teen verlaagde dosisse toedien nie. 8. Moet nie Azox Forte 50 WG op druiwestokkies in kwekerie toedien nie.	