



# DYNAMO 800 SC

Reg. No. L9619, Act No. 36 of 1947 | Reg. Nr. L9619, Wet Nr. 36 van 1947

A suspension concentrate herbicide with long residual action for the control of annual weeds as listed in various crops as listed as well as for industrial use.

## HERBICIDE GROUP CODE C2

### ACTIVE INGREDIENT:

Diuron (substituted urea) 800 g/l

Manufactured for and Registered by | Vervaardig vir en Geregistreer deur:

### Sharda International Africa (PTY) LTD

Reg. No./Reg. Nr. 2010/002268/07

P. O. Box/Posbus 82021, Southdale, 2135

Tel: 031-764 3011

Tel: 087-822 2397

'n Suspensie konsentraat onkruidoder met lang nawerking vir die beheer van eenjarige onkruid soos gelys in verskeie gewasse soos gelys, sowel as vir industriële gebruik.

## ONKRUIDODER GROEPKODE C2

### AKTIEWE BESTANDDEEL:

Diuron (gesubstitueerde ureum) 800 g/l

BATCH NO. LOT NR.	
EXPIRY DATE VERVAL DATUM	

NET CONTENTS /  
NETTO INHOUD

20 l

UN No./VN Nr. 3082

© Sharda International Africa cc



CAUTION  
VERSIGTIG



**WARNINGS**

- Handle with care.
- Harmful if swallowed.
- Keep out of reach of children, uninformed persons and animals.
- May cause irritation of the eyes.
- Toxic to aquatic organisms, and may cause long term adverse effects in the aquatic environment.
- Store in the original container in a cool place under lock and key, away from food, feed, fertiliser and seed.
- **Re-entry interval:** Do not enter the treated field until the spray deposit has dried unless wearing protective clothing.

Although this herbicide has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of weeds against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

**PRECAUTIONS**

- Do not inhale fumes or spray mist.
- Wear protective clothing when handling the concentrate (face shield and gloves).
- Avoid contact with the skin and eyes.
- Wash skin with soap and water after use or in the case of accidental skin contact.
- Wash any contaminated clothing before re-use.
- Flush the eyes with clean water in the case of accidental eye contact.
- Do not eat, drink or smoke while mixing or applying or before washing hands and face and change of clothing.
- Prevent contamination of food, feedstuff, eating utensils and drinking water.
- Prevent the drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Do not apply near or onto desirable plants or trees, or on areas where their roots may extend, or in locations where the product may be washed or moved into contact with their roots.
- Rinse the empty container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsings to the contents of the spray tank before disposing of the container by perforation and flattening. Never use the container for any other purpose.
- Clean the applicator thoroughly after use and dispose of the wash water where it will not contaminate crops, grazing, rivers or dams.

**RESISTANCE WARNING:**

For resistance management "DYNAMO 800 SC" is a group code C2 herbicide. Any weed population may contain individuals naturally resistant to "DYNAMO 800 SC" and other group code C2 herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by "DYNAMO 800 SC" or any other group code C2 herbicide.

To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code. Alternate or tank mix with registered products from different herbicide group codes.
  - Integrate other control methods (chemical, cultural, biological) into weed control programmes.
- For specific information on resistance management contact the registration holder of this product.

**USE RESTRICTIONS:**

- Do not apply more than 12.0 l "DYNAMO 800 SC" per hectare per year on any crop.

**DIRECTIONS FOR USE: Use only as directed****COMPATIBILITY:**

"DYNAMO 800 SC" is compatible with "SHARDA ACETOCHLOR 900 EC", "SHARMIN 480 SL", "SHARMIN 720 SL", "BUZZIN 480 SC", "SHARDA HEXAZINONE 240 SL", "SHARDA HEXAZINONE 750 WG" and "SHARDA PARAQUAT 200 SL". Always strictly adhere to the mixing instructions, and always read and follow this label and the label for the product with which "DYNAMO 800 SC" is mixed. Water quality can affect compatibility. It is recommended that a compatibility test be carried out prior to use.

**MIXING INSTRUCTIONS:**

- Half fill the spray tank with water. Mix the required amount of "DYNAMO 800 SC" with clean water before adding it to the spray tank. Top up with clean water. Maintain agitation of the spray mixture during mixing and spraying to ensure thorough mixing.
- When spraying a tank mix, add the required amount of "DYNAMO 800 SC" to the spray tank first as described above. Add more water to the spray tank before adding the other tank mix product in a diluted form. Maintain agitation of the spray mixture during mixing and spraying to ensure thorough mixing.
- Do not allow any spray mixture to stand for an extended length of time, e.g. overnight.

**APPLICATION INSTRUCTIONS:**

- Apply in 200 – 400 l water/ha.
- Accurately calibrate the spray equipment prior to application.

**GENERAL:****Pre-emerge application:**

"DYNAMO 800 SC" should preferably be applied onto moist, cleanly cultivated, clod free soil. Rain or irrigation is required within one week after application for optimal results.

**Post-emerge application:**

Apply "DYNAMO 800 SC" to broadleaf weeds up to the 4 leaf stage and to grasses before tillering. "DYNAMO 800 SC" can be applied in combination with other registered herbicides, however the weeds should be actively growing and take into consideration the growth stage for optimal results.

**RECOMMENDED DOSAGE RATES:**  
**AGRICULTURE**

CROP	DOSAGE RATE (l/ha)	REMARKS
<b>AVOCADOS, BANANAS AND MANGOES</b> (established orchards only)	4.0 l/ha	Apply during the growing season onto cleanly cultivated, moist soil. Repeat 2 to 3 months later if necessary.
<b>PINEAPPLES</b> (post-plant or on established plants)	4.0 l/ha	Apply onto cleanly cultivated, moist soil. It is preferable to apply 2 applications of 2.0l/ha (with an interval of 4 months) than to apply a single application of 4.0 l/ha.
<b>MACADAMIA AND PECAN NUTS</b> (established trees)	2.0 l/ha	Apply during spring onto cleanly cultivated soil. Repeat application 2 to 3 months later.
<b>SUGARCANE</b> (plant and ratoon)	3.0 – 4.0 l/ha	<b>Use of "DYNAMO 800 SC" only:</b> Apply in 200 to 400 l water per hectare.  The higher dosage rate is recommended where severe weed infestation is expected or on heavier soils where longer residual activity is required.
<b>Pre-emergence</b>		
<b>Pre- and early post-emergence</b>	"DYNAMO 800 SC" 2.5 – 3 l/ha <b>AND</b> 2.5 l/ha "SHARDA ACETOCHLOR 900 EC" <b>OR</b> 6.2 l/ha "SHARMIN 480 SL" <b>OR</b> 4.13 l/ha "SHARMIN 720 SL" <b>OR</b> 4 l/ha "BUZZIN 480 SC" <b>OR</b> 1.5 l/ha "SHARDA PARAQUAT 200 SL"	<b>Use of "DYNAMO 800 SC" in combination with other herbicides:</b> Apply post-emerge treatments as a directed inter-row spray. For improved weed control, select a tank mix product according to the problem weeds.
(ratoon sugarcane only)	"DYNAMO 800 SC" 1.0 – 2.0 l/ha <b>AND</b> 1.25 l/ha "SHARDA HEXAZINONE 240 SL" <b>OR</b> 400 g/ha "SHARDA HEXAZINONE 750 WG"	Do not apply to plant sugarcane. Apply as a directed inter-row spray.
<b>Pre- and early post-emergence</b>		

**RECOMMENDED DOSAGE RATES:**  
**INDUSTRIAL USE**

SITE	DOSAGE RATE (l/ha)	SPOT TREATMENT	REMARKS
<b>Industrial sites, and along roads and railway tracks</b>	10.0	50 ml per 50 m <sup>2</sup>	Can be applied as a pre-emergence application ("DYNAMO 800 SC" only), or as an early post-emerge application in a tank mix with "SHARDA PARAQUAT 200 SL".

**WEEDS CONTROLLED:**

Billygoat weed	<i>Ageratum conyzoides</i>	Large thorn apple	<i>Datura ferox</i>
Perennial pigweed	<i>Amaranthus deflexus</i>	Crabfinger grass	<i>Digitaria sanguinalis</i>
Cape pigweed	<i>Amaranthus hybridus</i>	Goose grass	<i>Eleusine indica</i>
Thorny pigweed	<i>Amaranthus spinosus</i>	Painted milkweed	<i>Euphorbia geniculata</i> ***
Red pigweed	<i>Amaranthus thunbergii</i>	Painted milkweed	<i>Euphorbia heterophylla</i> ***
Spanish blackjack	<i>Bidens bipinnata</i>	Red euphorbia	<i>Euphorbia hirta</i> ***
Cosmos	<i>Bidens formosa</i>	Gallant soldier	<i>Galinsoga parviflora</i>
Common blackjack	<i>Bidens pilosa</i>	Bladder weed	<i>Hibiscus trionum</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>	Common buffalo grass	<i>Megathyrus maximus</i> (= <i>Panicum maximum</i> )**
White goosefoot	<i>Chenopodium album</i>	Sweet buffalo grass	<i>Panicum schinzii</i> **
Green goosefoot	<i>Chenopodium carinatum</i>	Common purslane	<i>Portulaca oleracea</i>
Wandering Jew	<i>Commelina benghalensis</i> **	Tropical richardia	<i>Richardia brasiliensis</i>
Field bindweed	<i>Convolvulus arvensis</i>	Dwarf marigold	<i>Schkuhria pinnata</i>
Flax-leaf fleabane	<i>Coryza bonariensis</i>	Bur bristle grass	<i>Setaria verticillata</i>
Canadian fleabane	<i>Coryza canadensis</i>	Tall khakiweed	<i>Tagetes minuta</i>
Yellow nutsedge	<i>Cyperus esculentus</i> **		

\*\*For improved weed control in sugarcane, "DYNAMO 800 SC" can be applied in a tank mix with "SHARDA ACETOCHLOR 900 EC", "SHARMIN 480 SL", "SHARMIN 720 SL", "BUZZIN 480 SC", "SHARDA HEXAZINONE 240 SL", "SHARDA HEXAZINONE 750 WG" or "SHARDA PARAQUAT 200 SL". Consult the label for the product with which "DYNAMO 800 SC" is to be applied, and use according to label instructions.

\*\*\*Suppression of weeds.

The following herbicides are registered by Sharda International Africa cc under Act No. 36 of 1947:  
 "SHARDA ACETOCHLOR 900 EC" (900 g Acetochlor/l, EC) – Reg. No. LXXXX  
 "SHARMIN 480 SL" (480 g 2,4-D Amine/l, SL) – Reg. No. LXXXX  
 "SHARMIN 720 SL" (720 g 2,4-D Amine/l, SL) – Reg. No. LXXXX  
 "BUZZIN 480 SC" (480 g Metribuzin/l, SC) – Reg. No. L9475  
 "SHARDA HEXAZINONE 240 SL" (240 g Hexazinone/l, SL) – Reg. No. LXXXX  
 "SHARDA HEXAZINONE 750 WG" (750 g Hexazinone/kg, WG) – Reg. No. L9487  
 "SHARDA PARAQUAT 200 SL" (200 g Paraquat/l, SL) – Reg. No. L9432

**WAARSKUWINGS**

- Hanteer versigtig.
- Skadelik indien ingesluk.
- Hou buite bereik van kinders, ongeligte persone en diere.
- Kan irritasie van die oë veroorsaak.
- Giftig vir waterorganismes en mag langtermyn negatiewe effekte in die akwatiese omgewing veroorsaak.
- Stoor agter slot en grendel, in die oorspronklike houder in 'n koel plek, weg van voedsel, voer, kunsmis en saad.
- Herbetreding:** Moet nie die behandelde area betree alvorens die spuitneerslag nie droog is nie, tensy beskermende klerasie gedra word.

Alhoewel hierdie onkruidodder omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van onkruid teen die betrokke middel, sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing, of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

**VOORSORGMATREËLS**

- Moet nie dampe of spuitnevel inasem nie.
- Dra beskermende klere (gesigskerm en handskoene) wanneer die konsentraat hanteer word.
- Vermid kontak met die vel en oë.
- Was met seep en water na gebruik of as die produk per ongeluk in kontak met die vel gekom het.
- Was besoedelde klere voordat dit weer gebruik word.
- In die geval van oogkontak, spoel oë met skoon water uit.
- Moenie eet, drink of rook tydens vermenging of toediening of voordat hande en gesig gewas en ander klere na gebruik aangetrek is nie.
- Voorkom die besoedeling van voedsel, voer, eetgerei en drinkwater.
- Voorkom die oorwaai van spuitmis na ander gewasse, weiding, riviere, damme of gebiede wat nie onder behandeling is nie.
- Moenie toedien naby gewenste plantegroei, of in die omgewing van sulke plante se wortels, of waar die onkruidodder na sulke plante se wortels kan loog of spoel nie.
- Spoel die leë houder driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10% van die houder. Gooi die spoelwater by die inhoud van die spuittenk voordat die houder vernietig word deur gate daarin te maak en plat te slaan. Moenie die houder vir enige ander doel gebruik nie.
- Maak spuittoerusting deeglik skoon na gebruik en gooi waswater waar dit nie gewasse, weiding, riviere of damme sal besoedel nie.

**WEERSTANDSWAARSKUWING:**

"DYNAMO 800 SC" is 'n groepkode C2 onkruidodder. Enige populasie van 'n spesifieke onkruid mag individueel insluit wat 'n natuurlike weerstand teen "DYNAMO 800 SC" of enige ander groepkode C2 onkruidodder het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruid populasie oorheers. Hierdie weerstandbiedende onkruid sal waarskynlik nie deur "DYNAMO 800 SC" of enige ander groepkode C2 onkruidodder beheer word nie.

Om weerstand teen onkruidodders te vertraag:

- Vermid die eksklusiewe herhaaldelike gebruik van onkruidodders met dieselfde groepkode. Wissel af met, of gebruik tenkmengsels van, geregistreerde produkte in verskillende onkruidodder-groepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in die onkruidodder programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

**GEBRUIKSBEPERKINGS:**

- Moet nie meer as 12.0 l "DYNAMO 800 SC" per hektaar per jaar op enige gewas toedien nie.

**GEBRUIKSANWYSINGS: Gebruik slegs soos aangedui****VEREENIGBAARHEID:**

"DYNAMO 800 SC" is verenigbaar met "SHARDA ACETOCHLOR 900 EC", "SHARMIN 480 SL", "SHARMIN 720 SL", "BUZZIN 480 SC", "SHARDA HEXAZINONE 240 SL", "SHARDA HEXAZINONE 750 WG" en "PARAQUAT 200 SL". Voldoen altyd streng aan die menginstruksies, en lees en volg altyd die aanduidings van die etiket, sowel as die etiket vir die produk waarmee "DYNAMO 800 SC" gemeng gaan word. Die kwaliteit van die water kan wel verenigbaarheid aftekeer. Dit word aanbeveel dat 'n verenigbaarheid toets uitgevoer word voor gebruik daarvan.

**MENGINSTRUKSIES:**

- Vul helfte van die spuittenk met water. Meng die benodigde hoeveelheid "DYNAMO 800 SC" met 'n bietjie skoon water voordat dit by by spuittenk bygevoeg word. Vul die tenk verder met skoon water. Dit is belangrik om die mengsel voortdurend te roer sodat die middel in suspensie bly.
- Wanneer 'n tenkmengsel gespuit word, voeg eerste die verlangde hoeveelheid "DYNAMO 800 SC" in die spuittenk soos hierbo beskryf. Voeg meer water in die spuittenk by voordat die ander verdunde produk bygevoeg word. Dit is belangrik om die mengsel voortdurend te roer sodat die middel in suspensie bly.
- Moet nie die mengsel laat staan nie, bv. oornag.

**TOEDIENINGSINSTRUKSIES:**

- Dien toe in 200 – 400 l water/ha.
- Maak seker dat die toerusting akkuraat gekalibreer is voor toediening.

**ALGEMEEN:****Vooropkoms toediening:**

"DYNAMO 800 SC" moet verkieslik toegedien word op klam, skoonbewerkte, kluitvrye grond. Reën of besproeiing word binne een week na toediening benodig vir optimale resultate.

**Naopkoms toediening:**

Dien "DYNAMO 800 SC" op breëblaaronkruid toe tot op die 4-blaarstadium en op grasse voor spruitvorming. "DYNAMO 800 SC" kan toegedien word in kombinasie met ander geregistreerde onkruidodders, maar die onkruid moet aktief besig wees om die groei en die groeistadium met in ag geneem word vir optimale resultate.

**AANBEVOLE DOSISSE: LANDBOU**

GEWAS	DOSIS (l/ha)	OPMERKINGS
<b>AVOKADO'S, PIESANGS EN MANGO'S</b> (slegs op gevestigde boorde)	4.0 l/ha	Dien toe tydens die groeiseisoen op skoonbewerkte, klam grond. Herhaal 2 tot 3 maande later indien nodig.
<b>PYNAPPELS</b> (direk nadat geplant is of op gevestigde plante)	4.0 l/ha	Dien toe op skoonbewerkte, klam grond. Dit is beter om twee opeenvolgende bespuitings teen 2.0 l/ha (met 'n interval van 4 maande) toe te dien, as 'n enkelbespuiting van 4.0 l/ha.
<b>MAKADAMIA EN PEKANNEUTE</b> (gevestigde bome)	2.0 l/ha	Dien toe in die lente op skoonbewerkte grond. Herhaal bespuiting na 2 tot 3 maande.
<b>SUIKERRIET</b> (plant- en opslagriet)	3.0 – 4.0 l/ha	<b>Gebruik van "DYNAMO 800 SC" alleenlik:</b> Dien toe in 200 tot 400 l water per hektaar.  <b>Vooropkoms</b> Die hoër dosis word aanbeveel waar ernstige onkruid infestasië verwag word of op swaarder gronde waar 'n langer nawerking verlang word.
<b>Voor- en vroeg naopkoms</b>	"DYNAMO 800 SC" 2.5 – 3 l/ha <b>EN</b> 2.5 l/ha "SHARDA ACETOCHLOR 900 EC" <b>OF</b> 6.2 l/ha "SHARMIN 480 SL" <b>OF</b> 4.13 l/ha "SHARMIN 720 SL" <b>OF</b> 4 l/ha "BUZZIN 480 SC" <b>OF</b> 1.5 l/ha "SHARDA PARAQUAT 200 SL"	<b>Gebruik van "DYNAMO 800 SC" in kombinasie met ander onkruidodders:</b> Vir naopkoms beheer, dien behandelings toe as 'n gerigte tussen-ry bespuiting. Vir verbeterde onkruidbeheer, kies 'n tenkmengsel produk volgens die probleem onkruid.
(slegs op opslagriet)	"DYNAMO 800 SC" 1.0 – 2.0 l/ha <b>EN</b> 1.25 l/ha "SHARDA HEXAZINONE 240 SL" <b>OF</b> 400 g/ha "SHARDA HEXAZINONE 750 WG"	Moet nie op planriet gebruik nie. Dien toe as 'n gerigte tussen-ry bespuiting.
<b>Voor- en vroeg naopkoms</b>	"DYNAMO 800 SC" 1.0 – 2.0 l/ha <b>EN</b> 1.25 l/ha "SHARDA HEXAZINONE 240 SL" <b>OF</b> 400 g/ha "SHARDA HEXAZINONE 750 WG"	

**AANBEVOLE DOSISSE: INDUSTRIËLE GEBRUIK**

PLEK VAN TOEDIENING	DOSIS (l/ha)	KOL BEHANDELING	OPMERKINGS
Op nywerheids-perselle en langs paaië en spoorlyne	10.0	50 ml per 50 m <sup>2</sup>	Kan toegedien word as 'n vooropkoms toediening ("DYNAMO 800 SC" alleenlik), of in 'n tenk mengsel met "SHARDA PARAQUAT 200 SL" (vroeg naopkoms).

**ONKRUIDE WAT BEHEER WORD:**

Bokkruid	<i>Ageratum conyzoides</i>	Kruisvingergras	<i>Digitaria sanguinalis</i>
Meerjarige misbredie	<i>Amaranthus deflexus</i>	Jongosgras	<i>Eleusine indica</i>
Kaapse misbredie	<i>Amaranthus hybridus</i>	Gekleurde melkkruid	<i>Euphorbia geniculata</i> ***
Doringmisbredie	<i>Amaranthus spinosus</i>	Gekleurde melkkruid	<i>Euphorbia heterophylla</i> ***
Roomisbredie	<i>Amaranthus thunbergii</i>	Roosmelkkruid	<i>Euphorbia hirta</i> ***
Spaanse knapsekêrel	<i>Bidens bipinnata</i>	Knopkruid	<i>Galinsoga parviflora</i>
Kosmos	<i>Bidens formosa</i>	Terblansbossie	<i>Hibiscus trionum</i>
Gewone knapsekêrel	<i>Bidens pilosa</i>	Gewone buffelsgras	<i>Megathyrus maximus</i> (= <i>Panicum maximum</i> )**
Herderstassie	<i>Capsella bursa-pastoris</i>	Soetbuffelsgras/ Landsbuffelsgras	<i>Panicum schinzii</i> **
Withondebossie	<i>Chenopodium album</i>	Gewone porslein	<i>Portulaca oleracea</i>
Groenhandebossie	<i>Chenopodium carinatum</i>	Tropiese Richardia	<i>Richardia brasiliensis</i>
Bengalse wandelende jood	<i>Commelina benghalensis</i> **	Kleinkakiebos	<i>Schuhria pinnata</i>
Akkerwinde	<i>Convolvulus arvensis</i>	Klitssetaria	<i>Setaria verticillata</i>
Kleinskraalhans	<i>Conyza bonariensis</i>	Smalblaartaaiman	<i>Tagetes minuta</i>
Kanadese skraalhans	<i>Conyza canadensis</i>		
Geeluintjie	<i>Cyperus esculentus</i> **		
Grootstinkblaar	<i>Datura ferox</i>		

\*\*\*Vir verbeterde onkruidbeheer in suikerriet, kan "DYNAMO 800 SC" toegedien word in 'n tenkmengsel saam met "SHARDA ACETOCHLOR 900 EC", "SHARMIN 480 SL", "SHARMIN 720 SL", "BUZZIN 480 SC", "SHARDA HEXAZINONE 240 SL", "SHARDA HEXAZINONE 750 WG" of "SHARDA PARAQUAT 200 SL". Raadpleeg die etiket vir die produk waarmee "DYNAMO 800 SC" toegedien moet word, en voldoen aan alle etiketaanwysings.

\*\*\*Onderdrukking van onkruid.

Die volgende onkruidodders is geregistreer deur Sharda International Africa cc onder Wet Nr. 36 van 1947:  
 "SHARDA ACETOCHLOR 900 EC" (900 g Asetochoor/l, EC) – Reg. Nr. LXXXX  
 "SHARMIN 480 SL" (480 g 2,4-D Amien/l, SL) – Reg. Nr. LXXXX  
 "SHARMIN 720 SL" (720 g 2,4-D Amien/l, SL) – Reg. Nr. LXXXX  
 "BUZZIN 480 SC" (480 g Metribusien/l, SC) – Reg. Nr. L9475  
 "SHARDA HEXAZINONE 240 SL" (240 g Heksasinoon/l, SL) – Reg. Nr. LXXXX  
 "SHARDA HEXAZINONE 750 WG" (750 g Heksasinoon/kg, WG) – Reg. Nr. L9487  
 "SHARDA PARAQUAT 200 SL" (200 g Parakwat/l, SL) – Reg. Nr. L9432