

SHARDAROACHGEL 35g

Reg. No. L10674, Act No. 36 of 1947

A non-volatile, non-repellent gel bait insecticide for the control of cockroaches in domestic and food storage premises, as well as in public hygiene.

IRAC INSECTICIDE GROUP CODE 4A

Active ingredient: Imidacloprid (Nicotinoid) 21.5g/kg

WARNINGS

• Handle with care. • Harmful when swallowed. • Harmful to fish and other animals. • Store in a cool place away from food and feedstuffs. • Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

- Avoid contact by mouth or with skin.
- Wear rubber gloves during application.
- Wash with soap and water immediately after use or after accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying or before washing hands and face.
- Apply only in area where it will be completely inaccessible to children and pets.
- Do not contaminate dams, drains, ponds, streams and other waterways with the product or with used containers.
- Do not apply to surfaces on which food or feed is stored, prepared or eaten.
- Protect food preparing equipment and eating utensils from contamination during application.
- Destroy the empty container and do not re-use for any other purpose.

DIRECTIONS FOR USE Use only as indicated.

Suitable for use in commercial and industrial premises, warehouses, food handling and processing establishments, market places, restaurants, hotels, hospitals, barracks, ships, offices, store rooms, schools and homes, etc.

SHARDA ROACH GEL is a cockroach insecticide with delayed action; therefore the results will not be visible

Reg. Nr. L10674, Wet Nr. 36 van 1947

'n Nie-vlugtige, nie-afweermiddel-gel-aas insekdoder vir die beheer van kakkerlakke in huishoudelike en voedselopbergingspersele, sowel as in openbare higiëne.

IRAC INSEKDODER GROEP CODE 4A

Aktiewe bestanddeel: Imidakloprid (Nikotinoïed) 21.5g/kg

immediately. A noticeable effect on the cockroach population may only be observed within a week. The area to undergo treatment should be examined to identify the centres of the cockroach infestation.

SHARDA ROACH GEL can then be applied by spot-treatment in places in and around cockroach harbours and where cockroaches have been observed or where they hide. The gel should be applied in dark areas and well out of sight. **SHARDA ROACH GEL** is dispensed in an injection tube.

APPLICATIONS RATES: Bait application should consist of 0.25-0.5 g/spot and 10-20 spots/10 m². A spot of 3 mm in diameter is equal to approximately 0.25 g of bait whereas a spot of 6 mm in diameter equals approximately 0.5 g of bait. The lower application rate can be used for low and moderate infestations or re-treatments, while the higher rate can be used for more severe infestations or a primary treatment. Rapid control is provided by a number of small placements rather than less large spots. The gel will remain pliable

usually for up to 20 weeks after application. It is advisable to inspect the gel after 4 weeks of treatment and replenish if necessary. Daylight conditions will reduce the longevity of the bait. **SHARDA ROACH GEL** must not be contaminated with other insecticides or placed on freshly sprayed surfaces.

Sharda International Africa (Pty) LTD

Reg. No./Reg. Nr. 2010/002268/07

P.O. Box/Posbus 82021, Southdale, 2135

Tel: 031-764 3011 | Tel: 087-822-2397



**CAUTION
VERSIGTIG**



Batch No.
Expired Date
UN No./VN Nr. 3077

Lot Nr.
Verval Datum